

Radiopharmaceuticals for SPECT Cancer Detection

V. I. Chernov^{1,2,a)}, A. A. Medvedeva^{1,2,b)}, R. V. Zelchan^{1,2,c)}, I. G. Sinilkin^{1,2,d)},
E. S. Stasyuk², L. A. Larionova², E. M. Slonimskaya¹, and E. L. Choyzonov¹

¹ Tomsk Cancer Research Institute, Kooperativny Street 5, Tomsk, 634050 Russia

² Tomsk Polytechnic University, Lenin Avenue 30, Tomsk, 634050 Russia

^{a)} Corresponding author: chernov@oncology.tomsk.ru

^{b)} tickayaAA@oncology.tomsk.ru

^{c)} r.zelchan@yandex.ru

^{d)} sinilkinig@oncology.tomsk.ru

Abstract. The purpose of the study was to assess the efficacy of single photon emission computed tomography (SPECT) with ¹⁹⁹Tl and ^{99m}Tc-MIBI in the detection of breast, laryngeal and hypopharyngeal cancers. A total of 220 patients were included into the study: 120 patients with breast lesions (100 patients with breast cancer and 20 patients with benign breast tumors) and 100 patients with laryngeal/hypopharyngeal diseases (80 patients with laryngeal/hypopharyngeal cancer and 20 patients with benign laryngeal/hypopharyngeal lesions). No abnormal ¹⁹⁹Tl uptake was seen in all patients with benign breast and laryngeal lesions, indicating a 100% specificity of ¹⁹⁹Tl SPECT. In the breast cancer patients, the increased ¹⁹⁹Tl uptake in the breast was visualized in 94.8% patients, ^{99m}Tc-MIBI—in 93.4% patients. The increased ¹⁹⁹Tl uptake in axillary lymph nodes was detected in 60% patients, and ^{99m}Tc-MIBI—in 93.1% patients. In patients with laryngeal/hypopharyngeal cancer, the sensitivity of SPECT with ¹⁹⁹Tl and ^{99m}Tc-MIBI was 95%. The ¹⁹⁹Tl SPECT sensitivity in identification of regional lymph node metastases in the patients with laryngeal/hypopharyngeal cancer was 75% and the ^{99m}Tc-MIBI SPECT sensitivity was 17%. The data obtained showed that SPECT with ¹⁹⁹Tl and ^{99m}Tc-MIBI can be used as one of the additional imaging methods in detection of tumors.

INTRODUCTION

According to the recent WHO data more than 10 million newly diagnosed cancer cases occur annually in the world. In many countries cancer ranks the second most common cause of death following cardiovascular diseases 0.

Nuclear medicine techniques have the advantage over X-ray or magnetic resonance imaging in terms of higher specificity with a similar high sensitivity for cancer detection [1–3]. Moreover, the use of nuclear medicine imaging in detection of laryngeal/hypopharyngeal tumors allows prevention of unnecessary diagnostic invasive procedures.

^{99m}Tc-methoxy-isobutyl-isonitrile (^{99m}Tc-MIBI) is the most widely used single photon emission computed tomography (SPECT) tracer in detecting malignant disease [4, 5]. However, this tracer with a relatively high sensitivity has rather a low specificity. Therefore, the search for radiopharmaceuticals characterized by a high specificity while maintaining adequate sensitivity is of great importance.

More than 30 years ago the diagnostic value of ²⁰¹Tl for detecting malignant tumors was studied. It was shown that ²⁰¹Tl, a biological potassium analogue, accumulated actively in tumors [5]. The findings from animal studies demonstrated that ²⁰¹Tl had higher tumor accumulation compared to other radiopharmaceuticals [6, 7]. However, a long half-life (72 hours) of ²⁰¹Tl limited its widespread use, since there was a risk of increased radiation dose to the patient's body and critical organs, especially during re-examinations, for example during chemotherapy.

It is proposed that ¹⁹⁹Tl is a more promising isotope than ²⁰¹Tl, which is currently used in nuclear medicine. Unlike ²⁰¹Tl, ¹⁹⁹Tl has a simpler and cheaper production method. There are many reports on the use of this isotope for myocardial perfusion scintigraphy. Cellular uptake mechanism of ¹⁹⁹Tl is the same as that of ²⁰¹Tl. However, because of its short half-life (7.4 hours), the radiation dose to the patient's whole body and critical organs is

significantly reduced. By using ^{199}Tl , the radiation dose to critical organs is 4–15 times lower than that by using ^{201}Tl . It allows ^{199}Tl scintigraphy to be performed 5 times a year for the same patient [8–11].

The aim of our study was to evaluate the efficacy of SPECT with ^{199}Tl in the detection of breast and laryngeal/hypopharyngeal carcinomas.

MATERIALS AND METHODS

A total of 220 patients were included into the study. There were 120 women with breast disease. Out of them, 39 patients with stage $\text{T}_{1-4}\text{N}_{0-2}\text{M}_0$ breast cancer and 20 patients with benign breast lesions were injected with ^{199}Tl , and 61 patients with stage $\text{T}_{1-4}\text{N}_{0-2}\text{M}_0$ breast cancer were injected with $^{99\text{m}}\text{Tc}$ -MIBI. There were 80 patients with stage $\text{T}_{1-3}\text{N}_{0-2}\text{M}_0$ laryngeal and hypopharyngeal cancers (including 20 patients with recurrent laryngeal/hypopharyngeal cancer) and 20 patients with benign diseases. Out of them, 50 patients were injected with ^{199}Tl and the other 50 patients were injected with $^{99\text{m}}\text{Tc}$ -MIBI. None of the patients at the time of the survey showed evidence of distant metastasis.

Besides conventional imaging methods such as computed tomography, ultrasound, fibrolaryngoscopy and biopsy, all patients underwent preoperative SPECT with ^{199}Tl or $^{99\text{m}}\text{Tc}$ -MIBI. The patients were subsequently operated on and surgical specimens were examined histologically and immunohistochemically. The patients with benign breast lesions underwent sectoral resection of the breast with subsequent histological examination.

Single photon emission computed tomography was performed using a double-head gamma-camera (E.CAM 180, Siemens) equipped with parallel high energy collimators for ^{199}Tl and parallel low energy collimators for $^{99\text{m}}\text{Tc}$ -MIBI. The injection of ^{199}Tl at the dose of 260 MBq or $^{99\text{m}}\text{Tc}$ -MIBI at the dose of 740 MBq was made intravenously. In the patients with breast disease the injection was made into the antecubital vein contralaterally to the breast lesion or in the pedal vein. The images of the breast were obtained with the patients lying in the supine position with the arms raised over the head and with the chest, both breasts, axilla and the myocardium, included into the field of view. For obtaining SPECT images in the patients with laryngeal/hypopharyngeal cancer, the patients were also placed in a supine position, and the detector's field of the view encompassed the head and neck region and trachea to the bifurcation. A total of 32 projection images were recorded into a 64×64 matrix (30 seconds per projection), without a magnification factor. The scan images were analyzed using the manufacturer software (e.soft, Siemens, Germany). Three-dimensional images of the chest, sagittal, transverse and coronal sections were obtained. Single photon emission computed tomography scans were visually assessed. The images of contralateral areas were compared and asymmetrically increased radiotracer uptake was considered pathological.

RESULTS

Single photon emission computed tomography revealed no abnormal ^{199}Tl uptake in the chest wall and breasts of patients with benign breast lesions. Despite a small group of patients ($n = 20$), the specificity of SPECT with ^{199}Tl was 100%.

In breast cancer patients, the increased ^{199}Tl uptake was visualized in 37 (94.8%) of the 39 patients. Out of 6 patients with tumors less than 10 mm 4 showed abnormal radiotracer uptake (Fig. 1). In one case, a large mass detected by mammography and ultrasound appeared as a cluster of microcalcifications. Carrying out SPECT with $^{99\text{m}}\text{Tc}$ -MIBI revealed abnormal formations in the projection of the mammary glands in 59 (93.4%) out of 61 patients (Fig. 2). Tumors greater than 10 mm were detected by ^{199}Tl and $^{99\text{m}}\text{Tc}$ -MIBI SPECT in 100% of cases.

It is well known that a secondary infiltrative ductal breast carcinoma can be difficult to diagnose. In our study, the accuracy in differentiating malignant from benign breast lesions by SPECT with ^{199}Tl was 100%, as well as with $^{99\text{m}}\text{Tc}$ -MIBI (Figs. 3, 4).




FIGURE 1. SPECT with ^{199}Tl for the patient with stage T2N0M0 cancer in the left breast




FIGURE 2. SPECT with ^{99m}Tc -MIBI for the patient with stage T1N0M0 cancer in the left breast




FIGURE 3. SPECT with ^{199}Tl for the patient with secondary infiltrative ductal breast carcinoma in the right breast




FIGURE 4. SPECT with ^{99m}Tc -MIBI for the patient with secondary infiltrative ductal breast carcinoma in the right breast




FIGURE 5. SPECT with ^{199}Tl for the patient with multicentric breast cancer in the right breast

Identification of multicentric breast cancer is an important factor in diagnosis of breast cancer. In our study, multicentric breast cancer was diagnosed in 66.6% of the patients during the study with ^{199}Tl and in 46.1% with ^{99m}Tc -MIBI (Figs. 5, 6).

As regard to lymph node involvement, the increased ^{199}Tl uptake in the axillary lymph nodes was observed in 15 out of 25 patients (60%) with histologically verified metastases (Fig. 7). Only 2 out of 5 patients had abnormal radiotracer uptake in the pectoral and subclavicular lymph nodes (Fig. 8). Thus, the ^{199}Tl SPECT sensitivity for the detection of metastatic regional lymph nodes in breast cancer patients was 60%.

It was possible to visualize the modified axillary lymph nodes in almost all the cases during the study with ^{99m}Tc -MIBI: in 27 (93.1%) out of 29 patients with verified metastasis pathological inclusion of the ^{99m}Tc -MIBI in axillary region was detected (Fig. 9). In addition, pathological changes in the projection of pectoral lymph nodes, subclavian and subscapularis groups were detected in 5 patients. This technique allowed us to differentiate between different levels of lymph nodes (Fig. 10). The sensitivity of the SPECT with ^{99m}Tc -MIBI in identifying metastatic regional lymph nodes was 93.1%. The specificity of SPECT with ^{199}Tl was 100%, with ^{99m}Tc -MIBI—80%.




FIGURE 6. SPECT with ^{99m}Tc -MIBI for the patient with multicentric breast cancer in the right breast




FIGURE 7. SPECT with ^{199}Tl in the patient with stage T2N1M0 cancer in the left breast. Axillary lymph node visualization




FIGURE 8. SPECT with ^{199}Tl in the patient with stage T2N3M0 cancer in the right breast. Visualization of pectoral and subclavicular lymph nodes




FIGURE 9. SPECT with ^{99m}Tc -MIBI in the patient with stage T2N1M0 cancer in the left breast. Axillary lymph node visualization

In laryngeal/hypopharyngeal cancer patients, both ^{199}Tl and ^{99m}Tc -MIBI abnormal uptake was detected in 95% of the patients with primary laryngeal/hypopharyngeal cancer (Figs. 11, 12). It should be noted that despite limitations for SPECT in visualization of tumors less than 10 mm. The minimum lesion size was 4 mm in the greatest dimension. In 20 out of 20 patients with recurrent laryngeal/hypopharyngeal cancer an abnormal uptake both ^{199}Tl and ^{99m}Tc -MIBI was detected; it was 100% sensitivity of SPECT with ^{199}Tl and ^{99m}Tc -MIBI in the diagnosis of recurrent laryngeal/hypopharyngeal cancer.

The histological examination of specimens obtained by ultrasound-guided fine needle aspiration biopsy showed cervical lymph node metastases in 18.3% of cases. On SPECT the lymph nodes, affected by metastases, were visualized in the form of tracer hyperfixation focus of irregular shape with rough, indistinct boundaries, expressed by a nonuniform distribution of tracer, usually of low intensity.




FIGURE 10. SPECT with ^{99m}Tc -MIBI in the patient with stage T2N3M0 cancer in the right breast. Visualization of pectoral, subclavicular lymph nodes




FIGURE 11. SPECT with ^{199}Tl for the patient with primary laryngeal cancer




FIGURE 12. SPECT with ^{99m}Tc -MIBI for the patient with primary laryngeal cancer




FIGURE 13. SPECT with ^{199}Tl for the patient with primary laryngeal cancer and cervical lymph node metastases

Single photon emission computed tomography with ^{199}Tl could detect biopsy-proven lymph node metastases in 75% of cases (Fig. 13). When performing SPECT with ^{99m}Tc -MIBI we were able to identify the accumulation of the drug in the projection of lymph nodes, affected by metastases, in 2 (17%) out of 12 patients (Fig. 14).




FIGURE 14. SPECT with ^{99m}Tc -MIBI for the patient with primary laryngeal cancer and cervical lymph node metastases




FIGURE 15. Screens the zones physiological accumulation of ^{99m}Tc -MIBI in the projection of the thyroid gland and submandibular salivary glands



FIGURE 16. Physiological accumulation of ^{199}Tl

We assume that such a low information content of SPECT with ^{99m}Tc -MIBI in the detection of metastases in regional lymph nodes is primarily related to their small size and physiological, initially high accumulation of ^{99m}Tc -MIBI in the projection of the thyroid and submandibular salivary glands, which often screens the zones of the regional lymph collector (Fig. 15). This effect is less pronounced when using ^{199}Tl (Fig. 16).

DISCUSSION

Our first experience of using ^{199}Tl has clearly indicated that this radiopharmaceutical represents a useful diagnostic tool for the detection of breast and laryngeal/hypopharyngeal cancers. Single photon emission computed tomography with ^{199}Tl showed 100% specificity for detecting breast cancer, when compared to ^{99m}Tc -MIBI with reported specificity varying from 85% to 94% [5, 13].

With regard to the assessment of regional lymph node metastasis in patients with laryngeal/hypopharyngeal cancer, ^{199}Tl SPECT demonstrated high specificity, whereas ^{99m}Tc -MIBI failed to evaluate the status of regional lymph nodes due to its physiological high uptake in the thyroid gland.

Despite the abovementioned advantages, ^{199}Tl has also disadvantages. This cyclotron-produced radiopharmaceutical has a higher cost compared to ^{99m}Tc , the most widely used radioisotope in nuclear medicine. Moreover, our findings have demonstrated rather a low sensitivity in the detection of regional lymph node metastases in breast cancer patients, thus significantly impairing the diagnostic accuracy.

Anyway, ^{199}Tl is a promising radiopharmaceutical for the detection of malignant tumors; however, further studies are required to determine its role in the cancer diagnostic algorithm.

ACKNOWLEDGMENTS

The study reported in this article was conducted according to accepted ethical guidelines involving research in humans and/or animals and was approved by an appropriate institution or national research organization.

The study is compliant with the ethical standards as currently outlined in the Declaration of Helsinki.

All individual participants discussed in this study, or for whom any identifying information or image has been presented, have freely given their informed written consent for such information and/or image to be included in the published article.

REFERENCES

1. A. M. Granov, L. A. Tyutin, and M. S. Tolstanova, The application of positron emission tomography with 18-fluorodeoxyglucose in clinical oncology, *Problems Oncology* **49**(5), 563–573 (2003).
2. A. Aziz, R. Hashmi, and Y. Ogawa, Tc-99m-MIBI scintimammography, SPECT versus planar imaging, *J. Cancer Bioter. Radiopharm.* **14**(6), 495–500 (1999).
3. C. K. Kuhl, MRI of the breast cancer, *J. Eur. Radiol.* **10**, 46–58 (2000).
4. S. J. Kim, Y. T. Bae, J. S. Lee, I. J. Kim, and Y. K. Kim, Diagnostic performances of double-phase tc-99m MIBI scintimammography in patients with indeterminate ultrasound findings: visual and quantitative analyses, *J. Ann. Nucl. Med.* **21**(3), 145–150 (2007).
5. E. L. Choyzonov, R. V. Zelchan, A. A. Titskaya, I. G. Sinilkin, S. Y. Chijevskaya, and V. I. Chernov, SPECT with 99mTc-MIBI and 199Tl-chloride in the diagnosis of primary and recurrent malignant tumors of the larynx and hypopharynx, in *A Century of progress in head & neck cancer: 5th World Congress of IFHNOS & Annual Meeting of the AHNS, 2014, 26–30 July*, <http://ahns.jnabstracts.com/Detail.aspx?ID=56831>.
6. A. Sehweil, J. H. McKillop, and R. Milroy, 201Tl scintigraphy in the staging of lung cancer, breast cancer and lymphoma, *Nucl. Med. Commun.* **1**(4), 263–269 (1990).
7. R. P. Spencer, Tumour-seeking radiopharmaceuticals: nature and mechanisms, *Nuclear Med.*, **2**, 649–662 (1994).
8. Yu. B. Lishmanov, V. I. Chernov, and N. G. Krivonogov, Experimental perfusion scintigraphy using 199Tl-chloride, *Medical Radiology and Radiation Safety, Med. Radiolog. Radiatsion. Bezopasn.* **3**, 13–16 (1988).
9. V. Chernov, S. Triss, V. Skuridin, and Yu. Lishmanov, Thallium-199: a new radiopharmaceutical for myocardial perfusion imaging, *Int. J. Cardiac Imaging* **12**(2), 119–126 (1996).
10. Yu. B. Lishmanov, V. I. Chernov, S. V. Triss, and I. Yu. Mazurin, Thallium-199 myocardial scintigraphy, *Med. Radiolog.* **4**, 35–38 (1990).
11. A. A. Titskaya, V. I. Chernov, E. M. Slonimskaya, I. G. Sinilkin, and R. V. Zelchan, Mammoscintigraphy with 99mTc-MIBI in the diagnosis of breast cancer, *Siber. Med. J.* **25**(4-1), 92–95 (2010).