

XV International Scientific Conference “Chemistry and Chemical Engineering in XXI century”
dedicated to Professor L.P. Kulyov

Preparation and structure of double complex compounds [La(HMPA)₄(NO₃)₂][Cr(NH₃)₂(NCS)₄]

A.A. Bobrovnikova^a*, T.G. Cherkasova^a, A.G. Shirokolobova^a, L.M. Bolsunovskaya^b

^a T.F. Gorbachev Kuzbass State Technical University, Kemerovo, Russia

^b National Research Tomsk Polytechnic University, Tomsk, Russia

Abstract

This paper presents the results of chemical analysis, IR- spectroscopic analysis, thermal gravitational analysis, X-ray phase analysis, and X-ray structural analysis, conducted to determine the composition, structure and properties of the double complex salts - tetra(isotiocyanato)diaminechromates(III) of complex lanthanon(III) of ceric group with hexamethylphosphorotriamide (HMPA) and nitrate-groups as ligands.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

Peer-review under responsibility of Tomsk Polytechnic University

Keywords: double complex salts (DCS), rare-earth elements (REEs), tetra(isotiocyanato)diaminechromates(III) ammonium, hexamethylphosphorotriamide (HMPA), X-ray structural analysis.

1. Introduction

Rare-earth elements (REEs), their chemical compounds and products based on rare-earth metals are of particular interest and strategic importance for the modern industrial world. These elements are used in computer, telecommunication and optical technologies, in radio electronics, nuclear technology and other high-tech fields¹⁻⁸.

The priority task is to expand and deepen the knowledge of chemical and physicochemical properties of the REEs, as well as the synthesis of new compounds based on them.

* Corresponding author name. Tel.: +7-913-307-6638
E-mail address: bobrownickowa.al@yandex.ru

Lack of information about peculiarities of formation and properties of double complex salts (DCS) contained in crystal structures tetra (isotiocyanato)diaminechromates (III) - anion and complex cation of REEs, with HMPA, stipulated interest in the current research.

Tetra (isotiocyanato)diaminechromates(III) ammonium (Reineke salt) is an anionic complex compound which can be considered as a source of metaloligands, particularly, of chromium (III) ions, ammonia molecules and rhodanide groups⁹.

Hexamethylphosphorotriamide (HMPA) is widely used as a ligand of coordination compounds on the bases of various metals¹⁰. HMPA molecules can participate in coordination to a metal, protonated and hydrolyzed. It is possible to use compounds, containing an organic ligand, as precursors for making compositions of metal oxides, materials of the emissive layer¹⁰. The information of biological activity of lanthanoid compounds and the opportunities of their agricultural usage was obtained¹⁰.

DCS can be obtained in solid state by means of direct synthesis from aqueous solutions¹¹.

2. Content

The fine-crystalline precipitates were obtained; the composition of precipitants according to chemical analyses corresponds to empirical formula $C_{28}H_{78}CrN_{20}LnO_{10}P_4S_4$. These precipitates were obtained during the studies of the aqueous solutions of lanthanoid (III) nitrates $Ln(NO_3)_3 \cdot 6H_2O$ line «C.P.», where $Ln = La^{3+}, Ce^{3+}, Pr^{3+}, Nd^{3+}, Sm^{3+}, Eu^{3+}$ and Reineke salt $NH_4[Cr(NH_3)_2(NCS)_4] \cdot 0,5H_2O$, with molecular ratio 1:1, with later dropwise addition of organic ligand – HMPA, when pH is 6-7 and the solution's concentration is 0.05-0.5 mole/l.

The crystals are plate-shaped, with dimensions of 3-55 microns (Figure 1.). According to raster electron microscopy (SEM), X-ray structural analysis and X-ray fluorescence analysis it is possible to assume that the rare-earth elements are contained in the structure of compounds, do not form their own phase and are regularly distributed on the surface.

In order to determine the link between $Ln-O_{HMPA}$ the IR- spectroscopic analysis of the synthesized compounds is carried out. Group oscillations of double bonds ($P=O$) in free HMPA lie under $1350-1175\text{ cm}^{-1}$ ¹². These frequencies are recorded under 1302, 1262 cm^{-1} in spectra of the obtained compounds¹³.

The decrease in the oscillation frequency group of $P=O$ approximately by 60 cm^{-1} in comparison with the ligand spectrum indicates the electron density of the ligand molecule displacement to the central ion of the lanthanoid(III). Frequency of valence vibration (PNC) is shifted to higher values (992 cm^{-1}) in regard to oscillations in the free HMPA (982 cm^{-1}).




Fig. 1. SEM image of the surface of with tetra (isotiocyanato)diaminechromate(III) lanthanon(III) with HMPA

X-ray structural analysis of the solid product of nitrate lanthanon(III) with tetra(isotiocyanato)diaminechromates(III) ammonium in hydro- hexamethylphosphorotriamide solution showed the existence of a new compound of ionic type $[\text{La}(\text{HMPA})_4(\text{NO}_3)_2][\text{Cr}(\text{NH}_3)_2(\text{NCS})_4]$, (Fig.2)⁹. It was found that the symmetry of the tested sample is described by the monoclinic space group $P2_1/n$, $a=15.0360(3)$ Å, $b=15.1214(3)$ Å, $c=26.7529(7)$ Å, $\beta=100.6610(10)^\circ$, $V=5977.7(9)$ Å³, $Z=2$, $\rho=1.442$ g/cm³.




Fig. 2. Structure of compound $[\text{La}(\text{HMPA})_4(\text{NO}_3)_2][\text{Cr}(\text{NH}_3)_2(\text{NCS})_4]$, hydrogen atoms are not shown




Fig. 3. Crystalline packing in $[\text{La}(\text{HMPA})_4(\text{NO}_3)_2][\text{Cr}(\text{NH}_3)_2(\text{NCS})_4]$

Ion La^{3+} coordinates 8 oxygen atoms – four atoms from the two NO_3 -groups ($\text{La}-\text{O}$ 2.599(3)-2.635(3) Å), and four – from four monodentate molecules HMPA ($\text{La}-\text{O}$ 2.386 (3) -2.425 (3) Å). Bond lengths in the molecule HMPA are $\text{P}=\text{O}$ 1.498(3)-1.503(3) Å; $\text{P}-\text{N}$ 1.620(4)-1.653(4)^{10,14}.

In the crystal packing of compound $[\text{La}(\text{HMPA})_4(\text{NO}_3)_2][\text{Cr}(\text{NH}_3)_2(\text{NCS})_4]$ the system of hydrogen bonds occurs (Fig. 3). Anion $[\text{Cr}(\text{NH}_3)_2(\text{NCS})_4]^-$ hydrogen-bonds: NH_3 -groups with two molecules HMPA, one oxygen atom of one of the NO_3 -groups and two oxygen atoms of the second NO_3 -group.

The compounds thermolysis process involves several stages (Fig. 4). Exothermic peak in the temperature range 95-130°C corresponds to the decomposition of the complex with elimination and burning of ammonia molecules and the beginning of HMPA molecules removal. On IR-spectra of the decomposition products under 200°C, there is a decrease in intensity of the absorption bands of valence vibration of NH -groups ($3222, 3151 \text{ cm}^{-1}$) and group P-O ($1302, 1262 \text{ cm}^{-1}$).

The main mass loss occurs in the range of 200-500°C. On the curve of differential scanning calorimetry DSC the range is represented by two exo-effects with extreme values due to samples decomposition and products of decomposition burning.




Fig. 4. Thermogram of compound banding $[\text{La}(\text{HMPA})_4(\text{NO}_3)_2][\text{Cr}(\text{NH}_3)_2(\text{NCS})_4]$ on air

3. Conclusions

By varying such parameters as media acidity, concentration ranges and temperature compound compositions were obtained from aqueous solutions: $[\text{Ln}(\text{HMPA})_4(\text{NO}_3)_2][\text{Cr}(\text{NH}_3)_2(\text{NCS})_4]$, where $\text{Ln} = \text{La}^{3+}, \text{Ce}^{3+}, \text{Pr}^{3+}, \text{Nd}^{3+}, \text{Sm}^{3+}, \text{Eu}^{3+}$, HMPA – hexamethylphosphortriamide $((\text{CH}_3)_2\text{N})_3\text{PO}$. Isostructurality of obtained substances was proved due to the similarity of diffraction patterns, infrared spectra and picnometer density. X-ray structure analysis method for the direct determination of the structure $[\text{La}(\text{HMPA})_4(\text{NO}_3)_2][\text{Cr}(\text{NH}_3)_2(\text{NCS})_4]$ was successfully applied. X-ray diffraction characteristics and properties of the compounds may be used to conduct chemical analysis of crystals and to describe physicochemical properties, as well as their dependence on the various factors. It was determined that the thermal decomposition of DCS proceeds at relatively low temperatures without reversible color changes in the investigated temperature range.

The obtained results contribute to the fundamental knowledge of REE and also bring us closer to the controlled synthesis of new applicable materials.

Acknowledgements

The authors express deep gratitude to A.V. Virovets and E.V. Peresyphkina, the staff of A.V. Nikolaev Institute of Inorganic Chemistry, to E.S. Tatarinova, an assistant professor of chemistry, technology of inorganic substances and nanomaterials of KuzGSU, for research assistance, valuable advice and comments.

References

1. Cherkasova, T.G., Tatarinova, E.S., Mezentsev, K.V., Cherkasova, E.V., Isakova, I.V., Bobrovnikova, A.A. and others. Design, synthesis, and the prospect of new materials preparation on mono-and polynuclear based complexes with thiocyanate anions. *Materials of the II International Kazakhstan-Russian Conference on Chemistry and Chemical Technology*, 2012, 272-275.
2. Cherkasova, T.G., Tatarinova, E.S., Cherkasova, E.V., Isakova, I.V., Bobrovnikova, A.A., Tikhomirova A.V. and others. Double complex compounds - coordination precursors to create new materials. *Abstracts of the the IV International Conference of the D.I. Mendeleev Russian Chemical Society devoted to the 80th anniversary of P.D. Sarkisov birth*, 2012, 327-329.
3. Tang, Y., Zakharov, L.N., Kassel, W.S., Rheingold, A.L., Kemp R.A. Synthesis and structural characterization of solvated calcium amides containing bulky silylamide ligands. *Inorg. Chim. Acta*, 2005, **358**, 2014-2022.
4. Suss-Fink, G., Cuervo, L.G., Therrien, B., Stoeckli-Evans, H., Shulpin G.B. Mono and oligonuclear vanadium complexes as catalysts for alkane oxidation: synthesis, molecular structure, and catalytic potential. *Inorg. Chim. Acta*, 2004, **357**, 475-484.
5. Possibilities of application of rare earth elements to create structural materials for nuclear industry of the Ukraine. *Electronic resource*: http://vant.kipt.kharkov.ua/ARTICLE/VANT_2008_1/article_2008_1_195.pdf – Cap. from the screen.
6. Analytical chemistry section. *Electronic resource*: http://chem.usu.ru/win/conf/tesis99/anal/anal_thesis99.htm. – Cap. from the screen.
7. New fluorescent molecular materials based on coordination of rare earth elements compounds. *Electronic resource*: <http://rusnanotech09.rusnanoforum.ru/Public/LargeDocs/theses/rus/young/01/Shuvaev.pdf>. – Cap. from the screen.
8. Roussel, P., Boaretto, R., Kingsley, A.J., Alcock, N.W., Scot, P. Reactivity of a triamidoamine complex of trivalent uranium. *J. Chem. Soc., Dalton Trans*, 2002, 1423-1428.
9. Laser materials. – Dictionary of Scientific Terms – Medpulse.ru. *Electronic resource*: <http://www.medpulse.ru/encyclopedia/5517.html>. – Cap. from the screen.
10. Cherkasova, T.G. Cherkasova, E.V., Isakova, I.V., Kochnev, S.V., Bobrovnikova, A.A., Giniyatullina, Y.R.. Design, synthesis and properties of mono - and polynuclear thiocyanate complexes of transition metals and lanthanides. *Abstracts of the XIX Mendeleev Congress on General and Applied Chemistry*, 2011, 656.
11. Bobrovnikova A.A. Dissertation. Cand. chem. sciences. Kemerovo: KemSU, 2013. 120p.
12. Cherkasova, T.G., Isakova, I.V., Cherkasova, E.V., Tikhomirova, A.V., Kochnev, S.V., Bobrovnikova, A.A. and other. Thiocyanate and halide complexes of transition metals in the direct synthesis of double complex salts. *All-Russian scientific conference "Advances of synthesis complexing"*, 2011, 309.
13. Nakamoto, K. *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, M: World, 1991, 536.
14. Bobrovnikova, A.A. Synthesis and IR- spectroscopic analysis of reinekate of lanthanide of ceric group with hexamethylphosphorotriamide. *Polzunovskii Bulletin*, 2010, **3**, 71-73.
15. Bobrovnikova, A.A., Cherkasova, T.G., Tatarinova E.S. Crystal structures and physical and chemical properties of the lanthanide complexes with hexamethylphosphoramide, *Polzunovskii Bulletin*, 2013, **1**, 26-30.