



Article

Antioxidant Properties of New Phenothiazine Derivatives

Olesya Voronova ¹, Sergey Zhuravkov ², Elena Korotkova ¹, Anton Artamonov ³ and Evgenii Plotnikov ^{4,5,*} 

¹ School of Earth Sciences & Engineering, National Research Tomsk Polytechnic University, 30 Lenin Avenue, 634050 Tomsk, Russia; oaa@tpu.ru (O.V.); eikor@tpu.ru (E.K.)

² School of Nuclear Science & Engineering, National Research Tomsk Polytechnic University, 30 Lenin Avenue, 634050 Tomsk, Russia; zhuravkovsp@tpu.ru

³ Institute for Biomedical Problems, Russian Academy of Sciences, 76 A Khoroshevskoe, 123007 Moscow, Russia; anton.art.an@gmail.com

⁴ Research School of Chemistry & Applied Biomedical Sciences, National Research Tomsk Polytechnic University, 30 Lenin Avenue, 634050 Tomsk, Russia

⁵ Tomsk National Research Medical Center, Mental Health Research Institute, 4 Aleutskaya, 634014 Tomsk, Russia

* Correspondence: plotnikovev@tpu.ru

Abstract: Phenothiazine and its derivatives have a number of properties that contribute to their wider practical use in the production of biologically active substances, drugs, dyes, etc. Therefore, the synthesis and study of new compounds is of great relevance. The aim of this work was to investigate the antioxidant activity of a number of new phenothiazine derivatives. The patterns of electroreduction of oxygen and its radicals in the presence of phenothiazine derivatives in aqueous ethanol media were studied by voltammetry. The influence of various factors on antioxidant activity was considered by the methods of experiment planning. The optimal conditions for the manifestation of the antioxidant activity of phenothiazine derivatives have been found, which seems to be relevant since it opens up new possibilities for their further use as complex preparations with antioxidant activity, including in psychiatric practice.

Keywords: phenothiazine; phenothiazine derivatives; antioxidant activity; free radicals; voltammetry



Citation: Voronova, O.; Zhuravkov, S.; Korotkova, E.; Artamonov, A.; Plotnikov, E. Antioxidant Properties of New Phenothiazine Derivatives. *Antioxidants* **2022**, *11*, 1371. <https://doi.org/10.3390/antiox11071371>

Academic Editors: Guzel Ziyatdinova and Alla Ivanova

Received: 18 June 2022

Accepted: 13 July 2022

Published: 14 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Phenothiazine and its derivatives are one of the most important and promising groups of chemical compounds that are widely used in various fields of chemistry and medicine. These compounds are used in dyes, inhibitors of the oxidation and polymerization of monomers in liquid organic media, to stabilize polymers of various classes, and even photosensitizers [1–4]. They are characterized by a wide spectrum of action on the human body. The widespread use of phenothiazine derivatives is due to the high biological and pharmacological activity, which depends on the chemical structure of the compounds. Phenothiazine derivatives have revolutionized the treatment of mental illness, opening the era of psychopharmacology with the use of chlorpromazine [5]. Despite the emergence of new antipsychotics, phenothiazine derivatives are of great clinical importance in psychiatry to date. In wide medical practice, phenothiazine derivatives are also successfully used as antihistamines [6]. In addition, their antibacterial, antitumor, and antituberculosis effects are known [7–12]. The synthesis of new phenothiazine derivatives is still ongoing. It should be noted that an important nonspecific pathological process in a number of diseases is oxidative stress [13–17]. In all aspects of application, interaction with oxygen and its radicals becomes a key point for novel phenothiazine complexes. Previously, antioxidant properties have been shown in various phenothiazine derivatives [18,19]. Some phenothiazine derivatives exhibited reducing power potential to convert Fe(3+) to Fe(2+) and high ability to scavenge H₂O₂ free radical in vitro. These activities were comparable to ascorbic acid as a standard antioxidant [20]. For example, phenothiazine derivatives

inhibit the autoxidation of methyl linoleate [21]. Some authors have shown a prospective combination of antioxidant properties and biological activity [22]. Based on the literature data, we can predict high antioxidant properties in the compounds we have obtained. However, experimental confirmation of these properties and study of the mechanisms required in vitro tests. Further study will obviously include in vivo studies on biomodels of human diseases.

From this point of view, testing new compounds for their ability to neutralize free radicals is pathogenically justified. There are several methods for assessing the antioxidant and antiradical activity of substances [23–27]. We suppose that the methods mimicking the processes of oxygen reduction in living cells could most objectively reflect the behavior of an antioxidant in real objects [28–30]. We suggest here the optimal choice—the method of cathodic voltammetry, which is based on the electroreduction of oxygen, similar in stages to the processes in the mitochondria of a living cell. Electrochemical properties and patterns of electrode reactions for some phenothiazine derivatives have been established earlier [31]. Previously, we studied the antioxidant activity of some new compounds [32,33], and developed an effective method for assessing antioxidant properties [34]. Therefore, the main section of this work was the study of the antioxidant activity of new phenothiazine derivatives, which opens up new possibilities for their further use as antioxidants, including in the form of combined drugs.

The chemical structure of this compound is based on a heterocyclic system consisting of a thiazine heterocycle with two benzene rings (Figure 1). In this connection, they are good electron donors and easily enter into reactions with oxidation.




Figure 1. Structural formulas of phenothiazine and its derivatives: (a) phenothiazine (PT); (b) cis-10-propenylphenothiazine (cis-10-PPT); (c) 2-methyl-1-ethyl-1H-pyrido-[3,2,1-k,l]-phenothiazine. Pyridophenothiazine (PyrPT); (d) 2-methyl-1-ethyl-3-(10-phenothiazinyl)-2,3-dihydro-1H-pyrido-[3,2,1-k,l]phenothiazine. Propenylphenothiazine dimer (DPPT).

The main purpose of this work was to evaluate the antioxidant activity of phenothiazine and its derivatives using the method of cathodic voltammetry and to determine the optimal conditions for the measurement. The substances studied in this work potentially have high biological activity based on the chemical structure. However, this requires confirmation in experiments in vivo. This work allows us to consider these substances in a new way and significantly expand the potential of their use as antioxidants.

2. Materials and Methods

Phenothiazine powder (Sigma-Aldrich, Darmstadt, Germany) was used for the experiments. The rest of the investigated samples of substances were obtained for research by chemical synthesis according to the following method.

2.1. *Cis*-10-Propenylphenothiazine (*cis*-10-PPT)

To a suspension of 25 g (0.125 mol) of phenothiazine and 28 g of KOH in 250 cm³ of dry dimethyl sulfoxide at 80 °C, 25 cm³ (0.29 mol) of allyl bromide was added (Bp = 71–71.3 °C). The reaction solution was kept at the same temperature and stirred for 3 h. After complete conversion of phenothiazine, the reaction mixture was poured into 2000 cm³ of distilled water. The oily product was extracted from the water surface with benzene (three portions of 100 cm³); then, the benzene extract was washed and dried. The solvent was evaporated and the remnants of the reaction mass were distilled on the unit using a fore vacuum pump (at a temperature of 160 °C/1–2 mm Hg). The obtained 10-propenylphenothiazine (10-PPT) in the form of an oily liquid, collected in receiver flasks, was dissolved in portions with warm ethanol. After that, the flask was transferred to re-crystallization at a negative temperature. Then, after 5–10 h, the state of the mother liquor was checked—crystals should have fallen out of it. The precipitated crystals were filtered off on a Schott filter with a porosity of 160 pores. Then, the crystals were dried at room temperature. In total, 24.2 g (80%) of *Cis*-10-propenylphenothiazine (*Cis*-10-PPT) was received in the form of white needles with m.p. = 34–34.5 °C [35].

Gross formula of the obtained compound: C₁₅H₁₃NS.

Elemental analysis results:

Found, % (C 75.39; H 5.31; N 6.10; S 13.10).

Calculated, % (C 75.27; H 5.48; N 5.85; S 13.39).

In the IR spectrum of *Cis*-10-PPT, an intense band is observed in the region of 1657 cm^{−1}, and the band in the region of 945 cm^{−1} is attributed to the stretching and bending vibrations of the –C=C-bond, respectively (740 cm^{−1} (1,2-substituted benzene rings), 1600, 1500 cm^{−1} C=C aromatic rings). In addition, there is no absorption band of the free N-H bond in the IR spectra.

NMR spectrum: (CCl₄) δ ppm contains signals of the CH₃ methyl group in the region of 1.55 ppm (3H, *J* = 7 Hz), a weakly split doublet in the region of 6.16 ppm. =C-H (1H, *J* = 7 Hz) and a multiplet of protons of aromatic rings in the region of 6.5–7.1 ppm (8H).

2.2. 2-Methyl-1-ethyl-3-(10-phenothiazinyl)-2,3-dihydro-1H-pyrido-[3,2,1-*k,l*]phenothiazine Propenylphenothiazine Dimer (DPPT)

This substance was obtained by polymerization of 10-propenylphenothiazine at 20–25 °C in an organic solvent in the presence of a catalyst—boron trifluoride etherate in an amount of 3.6 × 10^{−3} mol per 1 mol of the starting monomer.

To do this, 2.4 g (0.01 M) of *cis*-10-propenylphenothiazine (*cis*-10-PPT) dissolved in 10 cm³ of an anhydrous organic solvent of ethyl acetate were loaded into the flask, 0.1 cm³ (0.00003 M) of BF₃·OEt₂ in ethyl acetate. The synthesis was carried out at 20–25 °C until complete conversion of *cis*-10-PPT. The progress of the reaction was monitored chromatographically (TLC, Silufol, Kavalier, Czech Republic, eluents: hexane:ether = 6:1). At the same time, 25–30 min after the addition of the catalyst to the reaction solution in the volume of the reaction mass, the appearance of crystals of the substance was observed, which then formed a precipitate on the walls of the flask. Then, within 1.5 h, a noticeable

increase in the volume of the precipitate was observed. At the end of the reaction, the catalyst was neutralized with a solution of KOH in methyl alcohol. The precipitate of the substance that accumulated during the synthesis was filtered off, washed, and dried. Then, 1.6 g (66.6%) of white small crystals were obtained with m.p. = 210–211°C, readily soluble in chloroform, benzene, poorly soluble in ethyl acetate, acetone, hexane [36].

Gross formula of the obtained substance: $C_{30}H_{26}N_2S_2$.

Elemental analysis results:

Found, % (C 76.70; H 5.69; N 5.43; S 12.99).

Calculated, % (C 75.31; H 5.43; N 5.85; S 13.38).

IR spectrum (Thermo Fisher Scientific (USA), FTIR spectrometer Nicolet 5700, KBr) 760 cm^{-1} (1,2-substituted benzene rings), $790, 740\text{ cm}^{-1}$ (1, 2, 3 substituted benzene rings), $1600, 1500\text{ cm}^{-1}$ C=C aromatic rings). In addition, there is no absorption band of the free N-H bond in the IR spectra.

NMR spectrum: ($CDCl_3$) δ ppm contains signals characteristic of the ethyl group: 1.02 (3H, t, $J = 6\text{ Hz}$), 1.28 (2H, d, $J = 7\text{ Hz}$), as well as signals in the regions 1.4–1.7 (2H, m), 2.4–2.85 (1H, m), 3.94–4.2 (1H, m), and 4.8–5.1 (1H, d, $J = 11\text{ Hz}$) and proton signals aromatic rings 6.5–7.2 (15H, m), $M^+ = 478$.

2.3. 2-Methyl-1-ethyl-1H-pyrido-[3,2,1-k,l]-phenothiazine

Pyridophenothiazine (PyrPT)

For synthesis, to a suspension of 1.0 g ($2.09 \times 10^{-3}\text{ mol}$) DPPT in 7.5 cm^3 of dioxane were added at 100°C in 0.24 cm^3 ($3.45 \times 10^{-3}\text{ mol}$) of acetyl chloride and 0.34 cm^3 ($4.18 \times 10^{-3}\text{ mol}$) of pyridine. The reaction solution was kept at the same temperature for 1 h. The reaction was carried out until complete conversion of the starting material. The progress of the reaction was monitored by thin layer chromatography. After completion of the reaction, the reaction mixture was cooled to room temperature. The precipitate formed was filtered off. The filtrate was poured into 50 cm^3 of water, the precipitate formed was filtered off. The precipitates were washed successively with 50 and 30 cm^3 of hexane. All washings and filtrates were combined. The organic phase was separated from the aqueous solution. The product was extracted from the aqueous solution with hexane. The extracts were dried. The solvent was evaporated, the precipitate was purified on a chromatographic column (silica gel sorbent, eluent—hexane). The resulting substance was recrystallized from ethanol. In total, 0.494 g (85%) of 2-methyl-1-ethyl-1H-pyrido[3,2,1-k,l]phenothiazine were obtained in the form of yellow crystals, m.p. = 70–71 °C [37].

Gross formula of the obtained compound: $C_{18}H_{17}NS$.

Elemental analysis results:

Found, %: C 77.41; H 6.13; N 5.00; S 11.46.

Calculated, %: C 77.34; H 6.09; N 5.01; S 11.48.

The IR spectrum of this compound indicates the absence of absorption of the free NH group and the presence of an absorption band of the olefin bond 1680 cm^{-1} , there are also bands: 740 and 790 cm^{-1} (1,2,3 substituted benzene ring), 755 cm^{-1} (1,2 substituted benzene ring).

NMR spectrum: ($CDCl_3$), δ , ppm: 1.0 (3H, t, $J = 6\text{ Hz}$, CH_2CH_3); 1.6 (2H, m, $CHCH_2Me$), 2.0 (3H, s, $=CCH_3$), 4.5 (1H, t $J = 6\text{ Hz}$, NCH), 6.1 (1H, s ArCH=) 6, 6–7.1 (7H, m HAr) phenothiazine ring protons.

Mass spectrum: $M^+ = 279$.

2.4. Antioxidant Activity Measurement


The measurements were carried out using an automated voltametric analyzer “TA-2” (LLC “Tomanalit”, Tomsk, Russia). The electrochemical cell consisted of a glass cup with a background electrolyte and holder with mercury-film indicator electrode, silver chloride reference electrode, and a silver chloride auxiliary electrode. Water–ethanol solution of 0.1 N $NaClO_4$ in various concentrations (10%, 30%, 50%, 70%, 85%) was used as a background solution.

The voltametric method for determining the antioxidant activity of phenothiazine and its derivatives consisted of recording voltammograms of cathodic oxygen reduction. We compared current pike with and without addition of test substances with different concentrations (0.01%, 0.1%, and 1.0%) to the background electrolyte solution in the potential range from 0.0 to -1.0 V to establish the relationship between the antioxidant activity of the studied substances and their concentration. The basic process here is oxygen electroreduction (O_2 ER); this process is identical to the oxygen reduction in the cells and tissues of the body. Notably, the mechanism of this process in monohydric alcohols is similar to that in water due to the identity of the structure of these media and the presence of hydrogen protons. The antioxidant activity of phenothiazine derivatives in such environment depends both on the concentration of the studied substances and on the water–ethanol ratio in the supporting electrolyte. The details of the voltametric method for measuring antioxidant activity is described previously in [34,38].

3. Results

3.1. Investigation of Oxygen Electroreduction in the Presence of Phenothiazine and Its Derivatives

The process of ER O_2 on the surface of a mercury-film electrode proceeds according to the scheme:



Since phenothiazine and its derivatives readily dissolve in aprotic solvents, we first considered their effect on the O_2 ER process in the aprotic solvent DMF. Thus, the effect of their interaction only with oxygen and the radical anion was evaluated (stage 1 according to the scheme). For a more correct interpretation of the results for living systems, we decided to use an ethanol solution, the mechanism of oxygen reduction that is similar due to the presence of hydrogen protons (stages 1–4)

In both cases, the substances exhibited antioxidant activity, reducing the current of ER O_2 , which indicates their interaction with oxygen and its radicals (Figure 2).

We studied the effect of the potential sweep rate on the current and potential of the oxygen reduction signal in the range from 20 to 200 mV/s to reveal the mechanism of O_2 ER in the presence of phenothiazine and its derivatives.

The peak current of the O_2 electroreduction in the presence of phenothiazine increases and the peak potential shifts to negative values with an increase in the potential scan rate (Figure 3). We studied dependences of the O_2 electrochemical reduction current on the sweep rate to the power ($W^{1/2}$) in the presence of the substances. It allows to reveal the reversibility of the electrochemical signal of O_2 reduction.

The current directly depends on the sweep rate to the power at $\frac{1}{2}$ degree for reversible and irreversible processes. However, a quasi-reversible process does not follow a linear relationship. The obtained dependences of the oxygen electroreduction current vs. the square root of the potential scan rate are nonlinear. This indicates the quasi-reversibility of the O_2 ER process in the presence of phenothiazine and its derivatives, as well as the O_2 ER process itself (Figure 4).

One of the criteria for the presence of a subsequent chemical reaction for quasi-reversible electrode processes is the presence of a linear dependence of the signal current peak potential on the logarithm of the potential scan rate to the power of $1/2$ ($\log W^{1/2}$).

However, nonlinear dependence was observed (Figure 5). Similar dependences were observed not only at 85% ethanol in the background electrolyte, but also for other more dilute ethanol concentrations.




Figure 2. Voltammogram of ER O₂ current 85% ethanol solution. 1—Background curve (0.1N NaClO₄); 2—In the presence of 0.1% phenothiazine.




Figure 3. Voltammograms of O₂ electroreduction on a mercury—film electrode versus the potential sweep rate in the presence of 0.1% phenothiazine in 85% ethanol solution.




Figure 4. Dependence of the O_2 electroreduction current on the RPE on the potential scan rate to the power of $1/2$ in the presence of 0.1% phenothiazine in 85% ethanol solution.




Figure 5. Dependence of the potential of the peak current of O_2 electroreduction on a mercury–film electrode on the potential scan rate to the power of $1/2$ in the presence of various concentrations of phenothiazine (0.01%, 0.1%, and 1%) in 85% ethanol solution.

The nonlinear dependence in various ethanol solutions confirms the occurrence of a quasi-reversible process of oxygen electroreduction. The mechanism of the influence of a substance on this process is similar at different compositions of the background electrolyte. There is also a potential shift towards negative values and peak stretching with an increase in the concentration of the studied substances in the background electrolyte solution (Figure 6). According to the literature data, this indirectly indicates the interaction of these substances by the CE mechanism with molecular oxygen dissolved in the electrolyte or the CEC mechanism with the preceding and subsequent chemical reactions of the interaction of the antioxidant with molecular oxygen and its reduction products (Figure 7).




Figure 6. O₂ electroreduction current voltammogram (mercury–film electrode), background curve (0.1N NaClO₄, 85% ethanol solution), and in the presence of various concentrations of phenothiazine (0.01%, 0.1%, and 1%).




Figure 7. Preceding and subsequent chemical reactions of the interaction of the antioxidant with molecular oxygen.

3.2. Determination of the Optimal Conditions for the Evaluation of Antioxidant Activity by the Experimental Design

Experimental planning methods were used to evaluate the optimal conditions for the antioxidant activity detection of the phenothiazine derivatives [28]. The antioxidant activity

of phenothiazine derivatives depends both on the concentration and on the water–ethanol ratio in the supporting electrolyte. Here, we used a full factorial experiment to build a mathematical model of the O₂ ER process in the presence of the substances under study, as well as assess its adequacy and assess the significance of the coefficients of the regression equation [39].

The variable factors were the concentration of the test substance (X₁) and the concentration of ethanol (X₂) in the supporting electrolyte. Since the oxygen concentration in the electrolyte with different ethanol concentrations is different, it is more correct to use the relative change in the O₂ ER current when the test substance is added. It directly corresponds to the antioxidant activity of the substance. Therefore, as a response function (Y), we used the relative change in the oxygen electroreduction current, expressed as a percentage according to the formula:

$$K^* = 100\% \left(1 - \frac{I_i}{I_o} \right) \cdot \frac{1}{t} \quad (5)$$

where I_i —value of limiting current ER O₂, mA; I_o —the value of the limiting current of ER O₂ in the absence of a substance in solution, mA; t —process time, min.

The main characteristics of the plan are presented in Table 1.

Table 1. Main characteristics of the plan N = 2².

Characteristic	X ₁ (C _{AO} , %)	X ₂ (C _{ethanol} , %)
Zero level	0.055	60
Variation interval	0.045	10
Upper level	0.1	70
Lower level	0.01	50

It was found that this model adequately describes the process for phenothiazine and its derivatives. All coefficients of the linear model are significant and have a plus sign, indicating that both factors have a positive effect on the response function. Moreover, the first factor (X₁)—the concentration of the test substance—has the greatest influence. The effect of the interaction of factors is not significant. Thus, the equations of the mathematical model are as follows (Table 2).

Table 2. The equations of the mathematical model.

Characteristic	Equations
PT	$Y = 0.923 + 0.168 X_1 + 0.098 X_2$
cis-10-PPT	$Y = 0.46 + 0.17 X_1 + 0.065 X_2$
PyrPT	$Y = 0.23 + 0.138 X_1 + 0.073 X_2$
DPPT	$Y = 0.129 + 0.027 X_1 + 0.022 X_2$

The response surface type “stationary elevation” is observed for all substances. It appears with a smooth increase in the response function with changing factors. The maximum is not detected here. The surfaces reach a plateau with an increase in the concentration of the studied substances. However, it was inappropriate to conduct studies at higher concentrations of phenothiazine derivatives, since the studies were based on a therapeutic dose (maximum daily dosage) of known drugs based on phenothiazine derivatives. Use of these drugs in large doses can lead to poisoning or death, and it should be taken into account when predicting medical use and selecting concentrations for biological tests. We used the kinetic criterion to assess the antioxidant activity of phenothiazine and its derivatives. Measurements were conducted taking into account

differences in the concentration of dissolved oxygen. This criterion reflects the amount of active oxygen radicals that reacted with the antioxidant per minute, K ($\mu\text{M}/\text{min}$):

$$K = \frac{C_{O_2}}{t} \left(1 - \frac{I_i}{I_0}\right) \quad (6)$$

where C_{O_2} —oxygen concentration in the initial solution without substance, μM ; I_i —value of the limiting current of O_2 electroreduction, mA ; I_0 —the value of the limiting current of electroreduction of O_2 in the absence of a substance in solution, mA ; t —process time, min .

Tables 3–6 present the results of a quantitative assessment of the antioxidant activity of substances with different concentrations (0.01%, 0.1%, and 1%) in water–ethanol solutions with different concentrations (30%, 50%, 70%, 85%).

Table 3. Values of the antioxidant activity of phenothiazine and its derivatives in 30% water–ethanol solution.

30% Water–Ethanol Solution	C1 = 0.01%		C2 = 0.1%		C3 = 1%	
	K, $\mu\text{M}/\text{min}$	Sr	K, $\mu\text{M}/\text{min}$	Sr	K, $\mu\text{M}/\text{min}$	Sr
PT	1.44	0.01	1.90	0.01	5.37	0.02
cis-10-PPT	6.48	0.01	12.80	0.03	19.43	0.04
PyrPT	6.64	0.02	10.59	0.03	13.11	0.03
DPPT	8.53	0.02	10.74	0.03	13.90	0.03

Table 4. Values of the antioxidant activity of phenothiazine and its derivatives in 50% water–ethanol solution.

50% Water–Ethanol Solution	C1 = 0.01%		C2 = 0.1%		C3 = 1%	
	K, $\mu\text{M}/\text{min}$	Sr	K, $\mu\text{M}/\text{min}$	Sr	K, $\mu\text{M}/\text{min}$	Sr
PT	4.73	0.02	9.65	0.02	17.11	0.03
cis-10-PPT	12.19	0.03	17.84	0.03	24.93	0.03
PyrPT	15.65	0.03	19.47	0.03	24.93	0.03
DPPT	15.29	0.03	23.84	0.04	44.77	0.04

Table 5. Values of the antioxidant activity of phenothiazine and its derivatives in 70% water–ethanol solution.

70% Water–Ethanol Solution	C1 = 0.01%		C2 = 0.1%		C3 = 1%	
	K, $\mu\text{M}/\text{min}$	Sr	K, $\mu\text{M}/\text{min}$	Sr	K, $\mu\text{M}/\text{min}$	Sr
PT	6.69	0.02	15.26	0.03	20.69	0.06
cis-10-PPT	17.56	0.03	21.32	0.03	31.35	0.07
PyrPT	19.86	0.03	26.13	0.03	29.47	0.07
DPPT	25.50	0.05	38.46	0.07	71.27	0.08

Table 6. Values of the antioxidant activity of phenothiazine and its derivatives in 85% water–ethanol solution.

85% Water–Ethanol Solution	C1 = 0.01%		C2 = 0.1%		C3 = 1%	
	K, $\mu\text{M}/\text{min}$	Sr	K, $\mu\text{M}/\text{min}$	Sr	K, $\mu\text{M}/\text{min}$	Sr
PT	17.78	0.05	23.03	0.06	24.85	0.06
cis-10-PPT	27.36	0.06	35.11	0.07	39.90	0.07
PyrPT	22.57	0.06	29.87	0.07	33.06	0.07
DPPT	39.90	0.07	65.44	0.08	82.76	0.08

Antioxidant activity increases in the row: phenothiazine < pyridophenothiazine < cis-10-propenylphenothiazine < propenylphenothiazine dimer. Antioxidant activity generally follows changes in the concentration of substances. There is a dose-dependent, but non-linear increase in activity with increasing concentration. In some cases, a change in the comparative activity in this series of compounds is noted at low concentrations. Here we revealed the highest antioxidant activity of the DPPT substance.

4. Discussion

The results confirmed that antioxidant activity increases with the concentration of the studied substances. However, it was found that this parameter is significantly affected by the water–ethanol ratio in the supporting electrolyte. It should be noted that the second factor (the ratio of solvents in the supporting electrolyte according to results of experiment planning) has the greatest influence on the parameter of antioxidant activity. The coefficients of antioxidant activity of the studied phenothiazine derivatives were calculated with correction of dissolved oxygen concentration. In the studied series, DPPT has the highest activity, and phenothiazine has the lowest activity. A possible explanation for the observed phenomenon is as follows. It is known that a feature of covalently bound sulfur in the structure of phenothiazine derivatives is high oxidizability. Oxidation processes can take place in several stages, and oxidation products are formed, including oxide and sulfur dioxide. We assume that the manifestation of propenylphenothiazine dimer with a significantly higher antioxidant activity compared to other derivatives is associated precisely with their molecular structure. As mentioned, the propenylphenothiazine dimer contains two fragments of the phenothiazine heterocycle, and pyridophenothiazine, which is its derivative, contains only one heterocycle. The observed dependence is associated with the chemical structure of thiazine heterocycle with two benzene rings. In this connection, these compounds are good electron donors and easily enter into reactions accompanied by oxidation. Notably, that the dual nature of phenothiazine exhibits both antioxidant and prooxidant properties, and this phenomenon is realized depending on the solvent or microenvironment in the biological object. Thus, in aqueous solutions, it exhibits mainly antioxidant properties, while in fats it behaves as a pro-oxidant with the formation of a phenothiazinyl radical [40]. In general, the antioxidant properties of phenothiazines are based on the chemical properties of the parent molecule, phenothiazine. However, the addition of functional groups significantly increases the antiradical properties of phenothiazine derivatives [20]. Possibly, antioxidant activity is enriched by the presence of electron donors such as sulfur as well as the richness of hydrogen in the additional benzene rings for substitution. However, some functional groups do not increase the activity of the molecule [41]. In addition, the presence of synergistic activity in combination with diphenylamine derivatives was noted, which is also an interesting phenomenon [42].

Evaluation of the influence of the phenothiazine derivatives on the reversibility of the O₂ ER process showed the following. The dependence of the oxygen electroreduction current on the sweep rate to the power of $\frac{1}{2}$ is nonlinear. This fact indicates the quasi-reversibility of the O₂ ER process on the mercury-film electrode in the presence of the studied substances.

Several possible mechanisms of interaction of the studied phenothiazine derivatives with oxygen and its active forms are suggested. The interaction of tested substances with molecular oxygen and oxygen reduction products goes by the chemical-electrochemical CE mechanism. This conclusion is indirectly confirmed by the absence of a linear dependence of the peak potential on the logarithm of the sweep rate to the power of $\frac{1}{2}$, as well as the shift of the potential towards negative values. At the same time, the CEC mechanism with the preceding and subsequent chemical reactions of the interaction of the antioxidant with molecular oxygen and the products of its reduction are not excluded.

To assess the potential use of these new antioxidants, a biological activity prediction was performed using the PASS online software. All tested substances are likely to have Glycosylphosphatidylinositol phospholipase D inhibitor properties. The probabilistic

indicator of activity was $P_a = 0.901$ (cis-10-PPT); $P_a = 0.915$ (Phenothiazine); $P_a = 0.571$ (Propenylphenothiazine dimer (DPPT)); and $P_a = 0.753$ (Pyridophenothiazine (PyrPT)). Here, the prediction of the biological activity of a chemical compound is based on the assumption that the activity is directly related to the structure. The probability of the presence of activity is expressed by the parameter P_a (probability “to be active”) that estimates the chance that the studied compound is belonging to the sub-class of active compounds. Accordingly, the closer calculated P_a value is to one, the higher the predicted probability of the compound having a certain activity. Thus, the high antioxidant activity that we identified can be manifested at the biological level through a specific inhibitory effect on phospholipase. It could significantly increase the antioxidant effect at the cellular level. It is known that glycosylphosphatidylinositol-specific phospholipase D may play an important role in inflammation, because it can hydrolyze the glycosylphosphatidylinositol anchors of several inflammatory membrane proteins. Therefore, the potential effect of the studied substances as inhibitors of glycosylphosphatidylinositol phospholipase and their high chemical antioxidant properties allows us to consider the studied substances as promising drugs for the correction of oxidative stress in various pathologies.

5. Conclusions

We proved high antioxidant activity of tested phenothiazine derivatives. The dependence of antioxidant activity on the molecular structure of the substance and the influence of various factors on their antioxidant activity was shown. It was revealed that the $ER O_2$ process is quasi-reversible in the presence of the studied substances. The interaction of these substances occurs presumably by the CE mechanism with molecular oxygen dissolved in the electrolyte or by the CEC mechanism with the preceding and subsequent chemical reactions of the interaction of the antioxidant with molecular oxygen and oxygen reduction products. The optimal conditions for the manifestation of the antioxidant activity of phenothiazine and its derivatives were found, which opens up new possibilities for their further use as antioxidants, including as combined drugs. Revealed antioxidant properties in combination of predicted biological activity make these substances promising for the correction of oxidative stress in the complex therapy of mental and cardiovascular diseases, where the role of oxidative damage has been proven.

Author Contributions: Conceptualization, E.P. and E.K. and O.V.; methodology, chemical synthesis S.Z.; software, A.A.; validation, O.V., S.Z. and E.K.; formal analysis, A.A.; investigation, O.V.; resources, O.V.; data curation, S.Z.; writing—original draft preparation, O.V.; writing—review and editing, A.A., O.V., E.P.; visualization, O.V.; supervision, E.K.; project administration, E.P.; funding acquisition, E.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by RFBR and Czech Science Foundation, grant # 19-53-26001; Russian State assignment “Science” # FSWW-2020-0022.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Available upon request.

Acknowledgments: This research was supported by TPU Development Program.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Xiau, Q. Pyridine-Embedded Phenothiazinium Dyes as Lysosome Targeted Photosensitizers for Highly Efficient Photodynamic Antitumor therapy. *J. Med. Chem.* **2020**, *63*, 4896–4907. [[CrossRef](#)] [[PubMed](#)]
2. Wainwright, M. Phenothiazinium photosensitisers, Part VI: Photobactericidal asymmetric derivatives. *Dyes Pigm.* **2009**, *82*, 387–391. [[CrossRef](#)]

3. Wilson, B. Synthesis and DNA photocleavage by mono- and bis-phenothiazinium-piperazine intercalators. *Tetrahedron* **2008**, *64*, 3429–3436. [\[CrossRef\]](#)
4. Ganeev, R. Peculiarities of the nonlinear optical absorption of Methylene blue and Thionine in different solvents. *Dyes Pigm.* **2018**, *149*, 236–241. [\[CrossRef\]](#)
5. López-Muñoz, F.; Alamo, C.; Cuenca, E.; Shen, W.W.; Clervoy, P.; Rubio, G. History of the discovery and clinical introduction of chlorpromazine. *Ann. Clin. Psychiatry* **2005**, *17*, 113–135. [\[CrossRef\]](#)
6. Cookson, J. Histamine in psychiatry: Promethazine as a sedative anticholinergic. *BJPsych Adv.* **2019**, *25*, 265–268. [\[CrossRef\]](#)
7. Grimsey, E.M.; Piddock, L.J.V. Do phenothiazines possess antimicrobial and efflux inhibitory properties? *FEMS Microb. Rev.* **2019**, *43*, 577–590. [\[CrossRef\]](#)
8. Motohashi, N.; Gollapudi, S.R.; Emrani, J.; Bhattiprolu, K.R. Antitumor properties of phenothiazines. *Cancer Investig.* **1991**, *9*, 305–319. [\[CrossRef\]](#)
9. Morak-Młodawska, B.; Jeleń, M.; Pluta, K. Phenothiazines Modified with the Pyridine Ring as Promising Anticancer Agents. *Life* **2021**, *11*, 206. [\[CrossRef\]](#)
10. Qi, L.; Ding, Y. Potential antitumor mechanisms of phenothiazine drugs. *Sci. China Life Sci.* **2013**, *56*, 1020–1027. [\[CrossRef\]](#)
11. Nizi, M.G.; Desantis, J.; Nakatani, Y.; Massari, S.; Mazzarella, M.A.; Shetye, G.; Sabatinia, S.; Barrecaa, M.L.; Manfronia, G.; Felicetti, T.; et al. Antitubercular polyhalogenated phenothiazines and phenoselenazine with reduced binding to CNS receptors. *Eur. J. Med. Chem.* **2020**, *201*, 112420. [\[CrossRef\]](#) [\[PubMed\]](#)
12. Warman, A.J.; Rito, T.S.; Fisher, N.E.; Moss, D.M.; Berry, N.G.; O'Neill, P.M.; Ward, S.A.; Biagini, G.A. Antitubercular pharmacodynamics of phenothiazines. *J. Antimicrob. Chemother.* **2013**, *68*, 869–880. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Plotnikov, E.; Korotkova, E.; Voronova, O.; Sazhina, N.; Petrova, E.; Artamonov, A.; Chernyavskaya, L.; Dorozhko, E. Comparative investigation of antioxidant activity of human serum blood by amperometric, voltammetric and chemiluminescent methods. *Arch. Med. Sci.* **2016**, *12*, 1071–1076. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Liguori, I.; Russo, G.; Curcio, F. Oxidative stress, aging, and diseases. *Clin. Interv. Aging* **2018**, *13*, 757–772. [\[CrossRef\]](#)
15. Rani, V.; Deep, G.; Singh, R.K.; Palle, K.; Yadav, U.C. Oxidative stress and metabolic disorders: Pathogenesis and therapeutic strategies. *Life Sci.* **2016**, *148*, 183–193. [\[CrossRef\]](#)
16. Cecerska-Heryć, E.; Polikowska, A.; Serwin, N.; Roszak, M.; Grygorcewicz, B.; Heryć, R.; Michalczyk, A.; Dołęgowska, B. Importance of oxidative stress in the pathogenesis, diagnosis, and monitoring of patients with neuropsychiatric disorders, a review. *Neurochem. Int.* **2022**, *153*, 105269. [\[CrossRef\]](#)
17. Forman, H.J.; Zhang, H. Targeting oxidative stress in disease: Promise and limitations of antioxidant therapy. *Nat. Rev. Drug Discov.* **2021**, *20*, 689–709. [\[CrossRef\]](#)
18. Murphy, C.M.; Ravner, H.; Smith, N.L. Mode of Action of Phenothiazine-Type Antioxidants. *Ind. Eng. Chem.* **1950**, *42*, 2479–2489. [\[CrossRef\]](#)
19. Dalla Tiezza, M.; Hamlin, T.A.; Bickelhaupt, F.M.; Orian, L. Radical Scavenging Potential of the Phenothiazine Scaffold: A Computational Analysis. *ChemMedChem* **2021**, *16*, 3763–3771. [\[CrossRef\]](#)
20. Engwa, G.A.; Ayuk, E.L.; Igbojekwe, B.U.; Unaegbu, M. Potential Antioxidant Activity of New Tetracyclic and Pentacyclic Nonlinear Phenothiazine Derivatives. *Biochem. Res. Int.* **2016**, *2016*, 9896575. [\[CrossRef\]](#)
21. Fukuzumi, K.; Ikeda, N.; Egawa, M. Phenothiazine derivatives as new antioxidants for the autoxidation of methyl linoleate and their reaction mechanisms. *J. Am. Oil Chem. Soc.* **1976**, *53*, 623–627. [\[CrossRef\]](#)
22. Al Zahrani, N.A.; El-Shishtawy, R.M.; Elaasser, M.M.; Asiri, A.M. Synthesis of Novel Chalcone-Based Phenothiazine Derivatives as Antioxidant and Anticancer Agents. *Molecules* **2020**, *25*, 4566. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Moharram, H.; Youssef, M. Methods for Determining the Antioxidant Activity: A Review. *Alex. J. Ed. Sci. Technol.* **2014**, *11*, 31–42.
24. Tirzitis, G.; Bartosz, G. Determination of antiradical and antioxidant activity: Basic principles and new insights. *Acta Biochim. Pol.* **2010**, *57*, 139–142. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Brand-Williams, W.; Cuvelier, M.E.; Berset, C. Use of a free radical method to evaluate antioxidant activity. *LWT- Food Sci. Technol.* **1995**, *28*, 25–30. [\[CrossRef\]](#)
26. Gulcin, İ. Antioxidants and antioxidant methods: An updated overview. *Arch. Toxicol.* **2020**, *94*, 651–715. [\[CrossRef\]](#)
27. Abeyrathne, E.D.N.S.; Nam, K.; Ahn, D.U. Analytical Methods for Lipid Oxidation and Antioxidant Capacity in Food Systems. *Antioxidants* **2021**, *10*, 1587. [\[CrossRef\]](#)
28. Sazhina, N. Determination of Antioxidant Activity of Various Bioantioxidants and Their Mixtures by the Amperometric Method. *Russ. J. Bioorg. Chem.* **2017**, *43*, 771–775. [\[CrossRef\]](#)
29. Siddeeg, A.; AlKehayez, N.M.; Abu-Hiamed, H.A.; Al-Sanea, E.A.; Al-Farga, A.M. Mode of action and determination of antioxidant activity in the dietary sources: An overview. *Saudi J. Biol. Sci.* **2021**, *28*, 1633–1644. [\[CrossRef\]](#)
30. Munteanu, I.G.; Apetrei, C. A Review on Electrochemical Sensors and Biosensors Used in Assessing Antioxidant Activity. *Antioxidants* **2022**, *11*, 584. [\[CrossRef\]](#)
31. Kuzin, Y.; Khadieva, A.; Padnya, P.; Khannanov, A.; Kutyreva, M.; Stoikov, I.; Evtugyn, G. Electrochemistry of new derivatives of phenothiazine: Electrode kinetics and electropolymerization conditions. *Electrochim. Acta* **2021**, *375*, 137985. [\[CrossRef\]](#)
32. Plotnikov, E.; Voronova, O.; Linert, W.; Martemianov, D.; Korotkova, E.; Dorozhko, E.; Astashkina, A.; Martemianova, I.; Ivanova, S.; Bokhan, N. Antioxidant and Immunotropic Properties of some Lithium Salts. *J. App. Pharm. Sci.* **2016**, *6*, 086–089. [\[CrossRef\]](#)

33. Plotnikov, E.; Korotkova, E.; Voronova, O.; Dorozhko, E.; Bohan, N.; Plotnikov, S. Lithium-based antioxidants: Electrochemical properties and influence on immune cells. *Physiol. Pharmacol.* **2015**, *19*, 107–113.
34. Korotkova, E.I.; Karbainov, Y.A.; Shevchuk, A.V. Study of antioxidant properties by voltammetry. *J. Electroanal. Chem.* **2002**, *518*, 56–60. [[CrossRef](#)]
35. Anfinogenov, V.; Napilkova, O.; Sirotkina, E.; Filimonov, V.D.; Ogorodnikov, V.D. Synthesis and cis-,trans-isomerization of 10-propenylphenothiazines. *Chem. Heterocycl. Compd.* **1986**, *1*, 121–124.
36. Anfinogenov, V.; Sirotkina, E.; Zhuravkov, S. 2-Methyl-1-ethyl-3-(10-phenothiazinyl)-2,3-dihydro-1H-pyrido-[3,2,1-k,l]-phenothiazine as a Stabilizer for Polyolefins. Patent RU #2105769, 27 February 1998.
37. Anfinogenov, V.; Sirotkina, E.; Zhuravkov, S. 1-ethyl-2-methyl-1n-pyrido[3,2,1-k,l]phenothiazine as an acceptor of alkyl radicals and a method for its preparation. Patent RU #2030410, 3 October 1995.
38. Miller, J.C.; Miller, J.N. *Statistics for Analytical Chemistry*, 6th ed.; Ellis Horwood PTR Prentice Hall: New York, NY, USA, 2010; p. 278.
39. Lazic, Z.R. *Design of Experiments in Chemical Engineering: A Practical Guide*; Wiley-VCH Verlag GmbH & Co. KGaA.: Weinheim, Germany, 2004; p. 610.
40. Iuga, C.; Campero, A.; Vivier-Bunge, A. Antioxidant vs. prooxidant action of phenothiazine in a biological environment in the presence of hydroxyl and hydroperoxyl radicals: A quantum chemistry study. *RSC Adv.* **2015**, *5*, 14678–14689. [[CrossRef](#)]
41. Morak-Młodawska, B.; Pluta, K.; Matralis, A.N.; Kourounakis, A.P. Antioxidant activity of newly synthesized 2,7-diazaphenothiazines. *Arch. Der Pharmazie.* **2010**, *343*, 268–273. [[CrossRef](#)]
42. Hui, W.; Zuwang, W.; Chaoliang, W.; Dong, S.; Fuli, W. Antioxidant Activity of 3,7-Di-iso-octyl-phenothiazine and Its Synergistic Effect with 4,4'-Di-iso-octyl-diphenylamine. *Tribol. Trans.* **2007**, *50*, 273–276. [[CrossRef](#)]